

Lymphatic drainage from the extracellular fluid of tumor tissue is impaired (leading to accumulation of macromolecules in this location)

- (1) A review of SMANCS. Maeda et al. J of protein Chem. 3, 181-193.
- (2) SMANCS and Polymer-Conjugated Macromolecular drugs. Maeda H. Adv. Drug Delivery Review, 1991, 6,181-202
- (3) Tumoritropic and Lymphotropic principles of Macromolecular Drugs, Maeda and Matsumura. Crit. Rev. Ther. Drug Carrier Systems, 1998, 6, 193-210
- (4) Use of Oily Contrast Medium for Selective Drug Targeting to Tumor. Enhanced Therapeutic Effect and X ray Image. Iwai et al. Cancer Research, 1984, 44, 2114-2121.
- (5) Tumor targeting by arterial administration of lipids. Iwai et al. Anticancer Research. 1987, 7, 321-328
- (6) Selective targeting of anticancer drug. Konno et al. Cancer, 1984, 54, 2367-2374
- (7) Image enhancement--and Selective Drug Targeting. Maki et al.Cancer, 1985, 56, 751-757
- (8) Effect of arterial administration of high molecular anticancer agents. Konno et al. European J. Cancer. Clinical Oncology, 1983, 19, 1053-1065
- (9) A new concept for macromolecular therapeutic. Mechanism of tumoritropic accumulation of proteins and anti-tumor agent smanc. Matsamuru and Maeda. Cancer Research. 1986, 46, 6387-6392
- (10) The interstitial water space of tumors. Guillino et al. Cancer Research. 1965, 25. 727-731

CERTIFICATE OF MAILING

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to Commissioner of Patents and Trademarks, Washington, D.C. 20231, on

JOHN McQUILLAN

(3)